



Med Instructions For Colonoscopy

DAY PRIOR TO PROCEDURE (Preparation Day)

Follow the clear liquid diet listed above and adjust your medication as follows:

If you take:	Morning	Lunch/Dinner	Bedtime
Actos (pioglitazone), Actoplus Met (pioglitazone & metformin), Avandia, Avandamet, Byetta, Farxiga, Glucophage (metformin), Invokana, Januvia, Janumet, Nesina, Onglyza, Tradjenta	Take your usual dose	Take your usual dose	
Amaryl (glimepiride), Avandaryl, DiaBeta, Glynase, Micronase (glyburide), Duetact (pioglitazone & glimepiride), Glucotrol, Glucotrol XL (glipizide), Glucovance (glyburide & metformin), Jardiance, Metaglip (glipizide & metformin HCl), Prandin (repaglinide), Starlix (nateglinide), Victoza	Do not take	Do not take	Do not take
Lantus Levemir	Take your usual dose		Take your usual dose
Humalog, Novolog, Regular Insulin or NPH Insulin If you use insulin sliding scale, you may continue using it.	Take ½ the usual amount if you take a fixed dose - OR - Cover your carbs with your usual carb ratio	Take ½ the usual amount if you take a fixed dose - OR - Cover your carbs with your usual carb ratio	
Premixed insulin; 75/25, Novolin 70/30 or Novolog Mix 70/30	Take ½ the usual dose at breakfast	Take ½ the usual dose at dinner	

DAY OF PROCEDURE:

Do not make up or "double up" on any missed medication after the procedure.

If you take:	Morning
Actos (pioglitazone), Actoplus Met (pioglitazone & metformin), Avandia, Avandamet, Byetta, Farxiga, Glucophage (metformin), Invokana, Januvia, Janumet, Nesina, Onglyza, Tradjenta	Take your usual dose
Amaryl (glimepiride), Avandaryl, DiaBeta, Glynase, Micronase (glyburide), Duetact (pioglitazone & glimepiride), Glucotrol, Glucotrol XL (glipizide), Glucovance (glyburide & metformin), Jardiance, Metaglip (glipizide & metformin HCl), Prandin (repaglinide), Starlix (nateglinide), Victoza	Do not take
Lantus, Levemir	Take ½ the usual dose
Humalog, Novolog, Regular Insulin or NPH Insulin. If you use insulin sliding scale, you may continue using it.	Do not take
Premixed insulin; 75/25, Novolin 70/30 or Novolog Mix 70/30	Do not take

- If you are taking any anti-platelet medications such as clopidogrel (Plavix) or ticlopidine (Ticlid) or blood-thinning medications, such as warfarin (Coumadin), please let us know. Depending on your clinical situation, we may advise you to stop these medications for 3 to 10 days before your procedure. We will need to consult your cardiologist or prescribing physician to make this recommendation. Please do NOT stop your medications before we are able to consult your cardiologist or prescribing physician and advise you.
- If you are taking a daily aspirin pill, please continue taking this medication.
- If you have diabetes, hypertension, or are taking a daily medication, please let us know so that we may advise you on whether or not you will need to stop your medication before your procedure. Please do NOT stop your medication before we are able to advise you.